arXiv:math/0004189v1 [math.NT] 7 Apr 2000

Elliptic Curves of Twin-Primes Over Gauss Field and Diophantine Equations [*†]

QIU Derong ( ) *and* ZHANG Xianke ( )

Tsinghua University, Department of Mathematical Sciences, Beijing 100084, P. R. China

Let p, q be twin prime numbers with $q - p = 2$. Consider the elliptic curves

$$E = E_\sigma : \quad y^2 = x(x + \sigma p)(x + \sigma q) . \qquad (\sigma = \pm 1) \tag{1}$$

$E = E_\sigma$ is also denoted as E_+ or E_- when $\sigma = +1$ or -1. Here the Mordell-Weil group and the rank of the elliptic curve E over the Gauss field $K = \mathbf{Q}(\sqrt{-1})$ (and over the rational field $\mathbf{Q}$) will be determined in several cases; and results on solutions of related Diophantine equations and simultaneous Pellian equations will be given. The arithmetic constructs over $\mathbf{Q}$ of the elliptic curve E have been studied in [1], the Selmer groups are determined, results on Mordell-Weil group, rank, Shafarevich-Tate group, and torsion subgroups are also obtained. Some results on torsion subgroups in [2] will be used here to determine $E(K)$.

Similarly to (1), some other special types of elliptic curves were studied by A. Bremner, J. Cassels, R. Strocker, J. Top, B. Buhler, B. Gross and D. Zagier (see [3-5]), *e.g.*, $y^2 = x(x^2 + p)$, $y^2 = (x + p)(x^2 + p^2)$, and $y^2 = 4x^3 - 28x + 25$. The last elliptic curve has rank 3 and is famous in solving the Gauss conjecture.

For any number field K, the Mordell-Weil theorem asserts that the $K-$rational points of E form a finitely generated abelian group (the Mordell-Weil group): $E(K) \cong E(K)_{tors} \oplus \mathbf{Z}^r$, where $E(K)_{tors}$ is the torsion subgroup, $r = \mathrm{rank}E(K)$ is the rank (see [6]).

Theorem 1. Let $E = E_\sigma$ be the elliptic curve in (1), $K = \mathbf{Q}(\sqrt{-1})$ the Gauss field.

(a) If $p \equiv 5 \pmod 8$, then

$$\mathrm{rank}E(K) = 0, \qquad E(K) \cong \mathbf{Z}/2\mathbf{Z} \oplus \mathbf{Z}/2\mathbf{Z}.$$

(b) If $p \equiv 3 \pmod 8$, $q = a^2 + b^2$, $(a + \epsilon)^2 + (b + \delta)^2 = c^2$ $(\epsilon, \delta = \pm 1,\ a, b, c \in \mathbf{Z})$, then

$$\mathrm{rank}E(K) = 1, \qquad E(K) \cong \mathbf{Z}/2\mathbf{Z} \oplus \mathbf{Z}/2\mathbf{Z} \oplus \mathbf{Z}.$$

(c) $\qquad \mathrm{rank}E(K) \leq 3.$

(d) If p, q are as in (b), then

$$\mathrm{rank}E_+(\mathbf{Q}) = 1, \qquad E_+(\mathbf{Q}) \cong \mathbf{Z}/2\mathbf{Z} \oplus \mathbf{Z}/2\mathbf{Z} \oplus \mathbf{Z}.$$

In [7], Euler considered the Diophantine equation

$$\begin{cases} X^2 + MY^2 = S^2 \\ X^2 + NY^2 = T^2 \end{cases}$$

for integers $M \neq N$, and studied the problem of classification of the couples (M, N) such that the above equation has non-trivial solutions, which is the famous"Euler Concordant Form problem". E. Bell, T. Ono, K. Ono studied this problem further in [8-10]. For twin prime numbers p, q we consider the following two Diophantine equations similar to the Euler's :

$$\begin{cases} X^2 - pY^2 = S^2 \\ X^2 - qY^2 = -T^2 \end{cases} \qquad (I)$$

$$\begin{cases} X^2 - pY^2 = 2S^2 \\ X^2 - qY^2 = -2T^2 \end{cases} \qquad (II)$$

If (X, Y, S, T) is a primary solution of (I) (or (II)) (i.e., an integer solution with $(X, Y) = 1$ and $XY \neq 0$), then by [1] we know that $(-X^2/Y^2,\ XST/Y^3)$ (or $(-X^2/Y^2,\ 2XST/Y^3)$) is a $\mathbf{Q}-$rational point of elliptic curve E_+ in (1). This fact leads us to the following theorem.

Theorem 2. If $p \equiv 5 \pmod 8$, then equations (I) and (II) have no primary solution. (Furthermore, the only solution of them is $(0, 0, 0, 0)$)

For elliptic curve $E = E_+$ in (1) and the Gauss field $K = \mathbf{Q}(\sqrt{-1})$, define

$$E(\mathbf{Q})^- = \{(x, y) \in E(\mathbf{Q}) \mid x < 0\},$$

$$n_{2K}^- = \frac{1}{2}\ \#(2E(K) \cap E(\mathbf{Q})^-),$$

and let $n(I)$ denote the cardinal of the set of positive primary solutions of equation (I).

Theorem 3 (3.1) If $n_{2K}^- \neq 0$, then equation (I) has a primary solution.

(3.2) We have $n_{2K}^- \leq n(I)$. In particular, if $n_{2K}^- = \infty$ then equation (I) has infinitely many primary solutions.

Finally, consider the simultaneous Pellian equations

$$\begin{cases} x^2 - py^2 = \sigma \\ z^2 - qy^2 = \sigma \end{cases} \qquad (\sigma = \pm 1) \qquad (III)$$

This kind of equations has been studied in [11-12] (but $\sigma = 1$ and $\{p, q\} = \{m, n\}$ are general integers). In particular, Rickert in [12] obtained: if $m = 2$, $n = 3$ then the equation has no nontrivial integer solution (Nontrivial means $y \neq 0$). We find that equation (III) has an interesting relation with elliptic curve E in (1): a nontrivial integer solution (a, b, c) of (III) gives a $\mathbf{Q}-$rational point $(1/b^2, ac/b^3)$ of E. We obtain a general result on equation (III).

Theorem 4. (a) If $p \equiv 5 \pmod 8$, then equation (III) has no nontrivial integer solution

References

[**1**] QIU Derong and ZHANG Xianke, Mordell-Weil Groups and Selmer Groups of Two Types of Elliptic Curves, to appear

[**2**] QIU Derong and ZHANG Xianke, Explicit determination of torsion subgroups of elliptic curves over multi-quadratic fields, Advances in Math., 28(1999), 475-478.

[**3**] A. Bremner and J. W. S. Cassels, On the equation $y^2 = x(x^2+p)$, Math. Comp. 42(1984), 257-264.

[**4**] R. J. Stroeker and J. Top, On the equation $y^2 = (x+p)(x^2+p^2)$, , Rocky Mountain J. of Math. 24(1994), 1135-1161.

[**5**] J. P. Buhler, B. H. Gross and D. B. Zagier, On the conjecture of Birch and Swinnerton-Dyer for an elliptic curve of rank 3, Math. Comp. 44(1985), 473-481.

[**6**] J. H. Silverman, The Arithmetic of Elliptic Curves, GTM 106, Springer-Verlag, 1986.

[**7**] L. Euler, De binis formulis speciei $xx + myy$ et $xx + nyy$ inter se concordibus et disconcordibus, Opera Omnia Series I, 5(1780), 48-60, Leipzig-Berlin-Zürich, 1944.

[**8**] E. T. Bell, The problems of congruent numbers and concordant forms, Proc. Amer. Acad. Sci., 33(1947), 326-328.

[**9**] T. Ono, Variations on a theme of Euler, Plenum, New York, 1994.

[**10**] K. Ono, Euler's condordant forms, Acta Arithmetica, LXXVIII. 2(1996), 101-123.

[**11**] H. P. Schlickewei, The number of subspaces occurring in the p-adic subspace theorem in Diophantine approximation, J, Reine Angew. Math., 406(1990), 44-108.

[**12**] J. Rickert, Simultaneous rational approximations and related Diophantine equations, Math. Proc. Cambridge Philos. Soc., 113(1993), 461-472.

TSINGHUA UNIVERITY
DEPARTMENT OF MATHEMATICAL SCIENCES
BEIJING 100084, P. R. CHINA
E-mail: xianke@tsinghua.edu.cn